## (19) World Intellectual Property Organization

International Bureau





(43) International Publication Date 30 September 2004 (30.09.2004)

### (10) International Publication Number WO 2004/083191 A1

- (51) International Patent Classification<sup>7</sup>: C07D 253/075, A61K 31/53
- (21) International Application Number:

PCT/IN2003/000057

- (22) International Filing Date: 17 March 2003 (17.03.2003)
- (25) Filing Language:

English

(26) Publication Language:

English

- (71) Applicant (for all designated States except US): HETERO DRUGS LIMITED [IN/IN]; Hetero House, 8-3-166/7/1, Erragadda, Hyderabad 500 018, Andhrapradesh (IN).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): PARTHASARADHI, Reddy, Bandi [IN/IN]; Hetero House, 8-3-166/7/1, Erragadda, Hyderabad 500 018, Andhrapradesh (IN). RATHNAKAR, Reddy, Kura [IN/IN]; Hetero Drugs Limited (R & D), Plot No. B-80 & 81, A.P.I.E., Balanagar, Hyderabad 500 018, Andhrapradesh (IN). RAJI, Reddy, Rapolu [IN/IN]; Hetero Drugs Limited (R & D), Plot No. B-80 & 81, A.P.I.E., Balanagar, Hyderabad 500 018, Andhrapradesh (IN). MURALIDHARA, Reddy, Dasari [IN/IN]; Hetero Drugs Limited (R & D), Plot No. B-80 & 81, A.P.I.E., Balanagar, Hyderabad 500 018, Andhrapradesh (IN). SUBASH CHANDER, Reddy, Kesireddy [IN/IN]; Hetero Drugs Limited (R & D), Plot No. B-80 & 81, A.P.I.E., Balanagar, Hyderabad 500 018, Andhrapradesh (IN).
- (74) Common Representative: RATHNAKAR, Reddy, Kura: Hetero Drugs Limited (R & D), Plot No. B-80 & 81. A.P.I.E., Balanagar, Hyderabad 500 018, Andhrapradesh (IN).

- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

#### Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- of inventorship (Rule 4.17(iv)) for US only

#### Published:

with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NOVEL CRYSTALLINE FORMS OF LAMOTRIGINE

(57) Abstract: The present invention relates to novel crystalline forms of lamotrigine, to processes for their preparation and pharmaceutical compositions containing them.



#### **NOVEL CRYSTALLINE FORMS OF LAMOTRIGINE**

#### FIELD OF THE INVENTION

The present invention relates to novel crystalline forms of lamotrigine, to processes for their preparation and pharmaceutical compositions containing them.

#### BACKGROUND OF THE INVENTION

10

20

25

Lamotrigine of formula (1):

$$CI$$
 $N$ 
 $N$ 
 $N$ 
 $N$ 
 $NH_2$ 

or 6-(2,3-Dichlorophenyl)-1,2,4-triazine-3,5-diamine is an anti-epileptic drug and its therapeutic uses are disclosed in U.S. Pat. No. 4,602,017.

Different synthetic methods of lamotrigine are described in WO 01/049669, US 6,111,101, US 6,333,198, US 5,912,345, EP 800521, US 4,602,017.

Various polymorphic forms are disclosed in WO 02/068398.

We have discovered three novel crystalline forms of lamotrigine. The novel forms have been found to be stable over the time and does not automatically convert into other crystalline forms of lamotrigine.

The novel forms of lamotrigine is, thus, suitable for pharmaceutical preparations.

Thus the object of the present invention is to provide stable novel crystalline forms of lamotrigine, to provide a processes for preparation of the



novel crystalline forms and to provide a pharmaceutical compositions comprising these novel crystalline forms.

#### SUMMARY OF THE INVENTION

5

10

15

20

30

According to one aspect of the present invention, there is provided a novel crystalline Form I of lamotrigine characterized by an x-ray powder diffraction pattern having peaks expressed as 20 at about 12.5, 13.9, 16.7, 18.0, 22.3, 23.6, 26.8, 27.9, 28.5, 28.9, 29.4, 31.7, 40.2, 42.3 degrees. Figure 1 shows typical Form I x-ray powder diffraction pattern.

According to another aspect of the present invention, there is provided a novel crystalline Form II of lamotrigine characterized by an x-ray powder diffraction pattern having peaks expressed as 20 at about 8.9, 11.2, 12.3, 13.2, 13.9, 17.0, 17.4, 18.0, 18.2, 18.7, 19.8, 21.4, 22.1, 22.7, 25.1, 25.4, 25.7, 26.4, 26.8, 27.1, 27.5, 28.3, 28.8, 29.2, 30.1, 30.9, 31.4, 32.9, 35.3, 35.7, 36.5 degrees. Figure 2 shows typical Form II x-ray powder diffraction pattern.

According to another aspect of the present invention, there is provided a novel crystalline Form III of lamotrigine characterized by an x-ray powder diffraction pattern having peaks expressed as 20 at about 9.9, 12.0, 13.3, 16.1, 17.0, 18.1, 18.7, 19.5, 24.4, 26.0, 26.3, 27.6, 28.1, 37.1 degrees. Figure 3 shows typical Form III x-ray powder diffraction pattern.

According to another aspect of the present invention there is provided a process for preparation of Form I lamotrigine comprising the steps of:

- a) dissolving lamotrigine in an ester;
- 25 b) maintaining at 15°C to 30°C for about 30 minutes to 2 hours;
  - c) filtering Form I lamotrigine.

The ester is selected from the group consisting of ethyl acetate, methyl acetate, ethyl formate, isopropyl acetate.

According to another aspect of the present invention there is provided a process for preparation of Form II lamotrigine comprising the steps of:

- a) dissolving lamotrigine in dioxane;
- b) maintaining at about 15°C to about 30°C for about 1 hour to 3 hours;
- c) filtering Form II lamotrigine.

According to another aspect of the present invention there is provided a process for preparation of Form III lamotrigine comprising the steps of:



- a) mixing lamotrigine, isopropyl acetate, chloroform and dimethyl formamide at about 60°C to about 70°C:
- b) filtering the Form III lamotrigine at about 20°C to about 30°C.
- Lamotrigine prepared by any of the known methods can be used in the above processes. Lamotrigine solvate or hydrate may also be used in the above processes.

According to another aspect of the present invention there is provided a pharmaceutical composition comprising Form I or Form III or Form III lamotrigine.

10

15

25

30

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a x-ray powder diffraction pattern of Form I lamotrigine. Figure 2 is a x-ray powder diffraction pattern of Form II lamotrigine. Figure 3 is a x-ray powder diffraction pattern of Form III lamotrigine. x-Ray powder diffraction spectrum was measured on a Siemens diffractometer.

#### **DETAILED DESCRIPTION OF THE INVENTION**

According to one aspect of the present invention, there is provided a novel crystalline Form I of lamotrigine characterized by an x-ray powder diffraction pattern having peaks expressed as 20 at about 12.5, 13.9, 16.7, 18.0, 22.3, 23.6, 26.8, 27.9, 28.5, 28.9, 29.4, 31.7, 40.2, 42.3 degrees. Figure 1 shows typical Form I x-ray powder diffraction pattern.

According to another aspect of the present invention, there is provided a process for preparation of Form I lamotrigine. Thus lamotrigine is dissolved in an ester. The ester is selected from the group consisting of ethyl acetate, methyl acetate, ethyl formate, isopropyl acetate. The Form I lamotrigine is maintained at about 15°C to about 30°C, preferably at about 20°C to about 25°C, for about 30 minutes to about 2 hours and filtered. Lamotrigine prepared by any of the known methods can be used in the process. Lamotrigine solvate or hydrate may also be used.

According to another aspect of the present invention, there is provided a novel crystalline Form II lamotrigine characterized by an x-ray powder diffraction

5

10

15

20

25

30



pattern having peaks expressed as 20 at about 8.9, 11.2, 12.3, 13.2, 13.9, 17.0, 17.4, 18.0, 18.2, 18.7, 19.8, 21.4, 22.1, 22.7, 25.1, 25.4, 25.7, 26.4, 26.8, 27.1, 27.5, 28.3, 28.8, 29.2, 30.1, 30.9, 31.4, 32.9, 35.3, 35.7, 36.5 degrees. Figure 2 shows typical Form II x-ray powder diffraction pattern.

According to another aspect of the present invention there is provided a process for preparation of Form II lamotrigine. Thus lamotrigine is dissolved in dioxane and maintained at about 15°C to about 30°C for about 1 hour to 3 hours. The separated Form II lamotrigine is filtered. Lamotrigine prepared by any of the known methods can be used in the process. Lamotrigine solvate or hydrate may also be used.

According to another aspect of the present invention, there is provided a novel crystalline Form III of lamotrigine characterized by an x-ray powder diffraction pattern having peaks expressed as 20 at about 9.9, 12.0, 13.3, 16.1, 17.0, 18.1, 18.7, 19.5, 24.4, 26.0, 26.3, 27.6, 28.1, 37.1 degrees. Figure 3 shows typical Form III x-ray powder diffraction pattern.

According to another aspect of the present invention there is provided a process for preparation of Form III lamotrigine. Thus lamotrigine isopropyl acetate, chloroform and dimethyl formamide are mixed and heated to about 60°C to about 70°C. The contents are maintained for about 30 minutes and cooled to about 20°C to about 30°C. The Form III lamotrigine is filtered. Lamotrigine prepared by any of the known methods can be used in the process. Lamotrigine solvate or hydrate may also be used.

According to another aspect of the present invention there is provided a pharmaceutical composition comprising Form I or Form II or Form III lamotrigine. The forms of lamotrigine may be formulated in a form suitable for oral administration or injection.

The following examples are given for the purpose of illustrating the present invention and should not be considered as limitations on the scope or spirit of the invention.

#### Example 1

Lamotrigine (10 gm) (obtained by the process described in example 1 of US 4,602,017) is mixed with ethyl acetate (100 ml) and maintained at about

5

25



70°C for 30 minutes. Then the contents are cooled to about 20°C. The solid is separated by filtration to give 9.0 gm of Form I lamotrigine.

#### Example 2

Lamotrigine (10 gm) (obtained by the process described in example 1 of US 4,602,017) is added to dioxane (100 ml), maintained at 50°C to 55°C for 30 minutes, cooled to 25°C and maintained at this temperature for 2 hours. The solid is separated by filtration to give 8.5 gm of Form II lamotrigine.

#### Example 3

Lamotrigine (10 gm) (obtained by the process described in example 1 of US 4,602,017) is added to isopropyl acetate (150 ml) and the contents are heated to about 65°C. Chloroform (50 ml) and dimethyl formamide (48 ml) are added at this temperature and stirred for 30 minutes. The contents are cooled to 25°C and filtered to give 9.5 gm of Form III lamotrigine.

#### Example 4

Example 1 is repeated using Form II lamotrigine instead of lamotrigine to give Form I lamotrigine.

#### Example 5

Example 1 is repeated using Form III lamotrigine instead of lamotrigine to give Form I lamotrigine.

20 Example 6

Example 2 is repeated using Form III lamotrigine instead of lamotrigine to give Form II lamotrigine.

#### Example 7

Example 2 is repeated using Form I lamotrigine instead of lamotrigine to give Form II lamotrigine.

#### Example 8

Example 3 is repeated using Form II lamotrigine instead of lamotrigine to give Form III lamotrigine.

#### Example 9


Example 3 is repeated using Form I lamotrigine instead of lamotrigine to give Form III lamotrigine.

20





#### We claim:


- 1. A crystalline lamotrigine Form I.
- 2. A crystalline lamotrigine, characterized by an x-ray powder diffraction pattern having peaks expressed as 20 at about 12.5, 13.9, 16.7, 18.0, 22.3, 23.6, 26.8, 27.9, 28.5, 28.9, 29.4, 31.7, 40.2, 42.3 degrees.
  - 3. A crystalline lamotrigine, characterized by an x-ray powder diffraction pattern as in figure 1.
- 4. A process for preparation of Form I lamotrigine of claim 1, comprising the steps of:
  - a) dissolving lamotrigine in an ester;
  - b) maintaining at 15°C to 30°C for about 30 minutes to 2 hours;
- c) filtering Form I lamotrigine;
  wherein ester is selected from the group consisting of ethyl acetate,
  methyl acetate, ethyl formate, isopropyl acetate.
  - 5. A crystalline lamotrigine Form II.
  - A crystalline lamotrigine, characterized by an x-ray powder diffraction pattern having peaks expressed as 20 at about 8.9, 11.2, 12.3, 13.2, 13.9, 17.0, 17.4, 18.0, 18.2, 18.7, 19.8, 21.4, 22.1, 22.7, 25.1, 25.4, 25.7, 26.4, 26.8, 27.1, 27.5, 28.3, 28.8, 29.2, 30.1, 30.9, 31.4, 32.9, 35.3, 35.7, 36.5 degrees.
  - 7. A crystalline lamotrigine, characterized by an x-ray powder diffraction pattern as in figure 2.
- 8. A process for preparation of Form II lamotrigine of claim 5, comprising the steps of:
  - a) dissolving lamotrigine in dioxane;
  - b) maintaining at about 15°C to about 30°C for about 1 hour to 3 hours;
  - c) filtering Form II lamotrigine.
  - 9. A crystalline lamotrigine Form III.
- 10. A crystalline lamotrigine, characterized by an x-ray powder diffraction pattern having peaks expressed as 2θ at about 9.9, 12.0, 13.3, 16.1, 17.0, 18.1, 18.7, 19.5, 24.4, 26.0, 26.3, 27.6, 28.1, 37.1 degrees.
  - 11. A crystalline lamotrigine, characterized by an x-ray powder diffraction pattern as in figure 3.



- 12. A process for preparation of Form III lamotrigine of claim 9, comprising the steps of:
- a) mixing lamotrigine, isopropyl acetate, chloroform and dimethyl formamide at about 60°C to about 70°C;
- 5 b) filtering the Form III lamotrigine at about 20°C to about 30°C.
  - 13. A pharmaceutical composition comprising a crystalline form of lamotrigine and a pharmaceutically acceptable carrier.
  - 14. A pharmaceutical composition of claim 13, wherein crystalline form of lamotrigine is Form I lamotrigine of claim 1.
- 15. A pharmaceutical composition of claim 13, wherein crystalline form of lamotrigine is Form II lamotrigine of claim 5.
  - 16. A pharmaceutical composition of claim 13, wherein crystalline form of lamotrigine is Form III lamotrigine of claim 9.









## INTERNATIONAL SEARCH REPORT

International application No. PCT/IN 03/00057-0

CLA	ASSIFICATION OF SUBJECT MATTER		
IPC <sup>7</sup> : C	07D 253/075, A61K 31/53		
	g to International Patent Classification (IPC) or to both na	ational classification and IPC	
	LDS SEARCHED documentation searched (classification system followed	hy classification symbols)	
	corp. A61K	by classification symbols	
	tation searched other than minimum documentation to the	e extent that such documents are included in	the fields searched
			a are rioras searoned
Electronic	data base consulted during the international search (name	ne of data base and, where practicable, search	ch terms used)
REGIS	TRY, CAPLUS		
C. DO	CUMENTS CONSIDERED TO BE RELEVANT		<del></del>
Category	Citation of document, with indication, where appropriate	e, of the relevant passages	Relevant to claim No.
Х	WO 2002/068398 A1 (Teva Pharmace Teva Pharmaceuticals USA, Inc.) 6 S cited in the application abstract, claim 1, fig. 1-18.		1-16
X	US 4602017 A (Baxter, Martin G.; Elpl A.; Sawyer, David A.; Wellcome Found 22 July 1986 (22.07.86); cited in the a abstract; example 1, col 6, lines 6-9.	1-16	
		<u> </u>	
* Specia "A" docum consid "E" earlier filing « "L" docum cited t specia "O" docum means "P" docum	nent which may throw doubts on priority claim(s) or which is o establish the publication date of another citation or other I reason (as specified) nent referring to an oral disclosure, use, exhibition or other	"T" later document published after the internation date and not in conflict with the application the principle or theory underlying the inversity document of particular relevance; the clair considered novel or cannot be considered to when the document is taken alone "Y" document of particular relevance; the clair considered to involve an inventive step who combined with one or more other such document obeing obvious to a person skilled in the arm. &" document member of the same patent family."	n but cited to understand nation ned invention cannot be o involve an inventive step ned invention cannot be nen the document is cuments, such combination
	ne actual completion of the international search	Date of mailing of the international search	report
	12 August 2003 (12.08.2003)	11 September 2003 (1	1.09.2003)
	I mailing adress of the ISA/AT	Authorized officer	
	in Patent Office	MÜLLER-HIEL	R.
	er Straße 87, A-1200 Vienna No. 1/53424/535	Talonhama No. 1/52/12/1/10/	
racsimile	No. 1/33424/333 T/ISA/210 (second sheet) (July 1998)	Telephone No. 1/53424/434	







International application No. PCT/IN 03/00057-0

Information on patent family members

	ent document cited In search report	Publication date		Patent memi	family ber(s)	Publicatio date
IL A	60201	31-07-1980	AR	A1	227521	15-11-198
IL Al	60201	31-05-1984	AT	A	2896/80	15-07-198
US A	4602017	22-07-1986	AT	В	370097	25-02-198
			ΑU	A1	58906/80	04-12-198
			AU	B2	530999	04-08-198
			BG	B2	60427	31-03-199
			CA	A1	1112643	17-11-198
			CA	A2	1133938	19-10-198
			CS	B2	234018	14-03-198
			CZ	A3	9103848	13-10-199
			ממ	С	151309	14-10-198
			DE	CO	3063084	16-06-198
			DE	C0	3071000	19-09-198
			DK	A	2338/80	02-12-198
			DK	В	153787	05-09-198
			DK	С	153787	16-01-198
			EP	Al	21121	07-01-198
	•		EP	Al	59987	15-09-198
			EP	B1	21121	11-05-198
			EP	B1	59987	14-08-198
			ES	A1	491998	16-05-198
			ES	A5	491998	15-06-198
			ES	A1	8104993	01-08-198
			FI	A	801758	02-12-198
			FI.	A	840888	06-03-198
			FI	A0	840888	06-03-198
		•	FI	В	67844	28-02-198
			FI	С	67844	10-06-198
			FI	В	73203	29-05-198
			FI	С	73203	10-09-198
			GR	Α	68380	28-12-198
			HU	В	182086	28-12-198
			IE	В	49823	25-12 <b>-</b> 198
			IT	AO	8048848	30-05-198
			IT	A	1147087	19-11-198
			JP JP	A2 A2	56025169	10-03-198
			JP	B4	61033163	17-02-198
			JP	B4 B4	1044179 1044706	26-09-1989
			LT	A3	2066	29-09-1989
			MY	A	62/85	15-06-1993
			N2	A	193890	31-12-1985
			NZ	A	198159	06-07-1984 09-11-1984
			PL	Al	224633	
			PL	B1	124029	13-02-1981 31-12-1982
			SU	A3	1055331	15-11-1983
			บร	A	4486354	04-12-1984
			YU	A	1456/80	28-02-1983
			ZA	A	8003250	27-01-1982
	-		ZW	A	129/80	06-01-1982
WO A1	02068398	06-09-2002	US	AA	03018030	23-01-2003
WO C2	02068398	21-11-2002				25 01-2003

# This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

□ BLACK BORDERS
□ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
□ FADED TEXT OR DRAWING
□ BLURRED OR ILLEGIBLE TEXT OR DRAWING
□ SKEWED/SLANTED IMAGES
□ COLOR OR BLACK AND WHITE PHOTOGRAPHS
□ GRAY SCALE DOCUMENTS
□ LINES OR MARKS ON ORIGINAL DOCUMENT
□ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

## IMAGES ARE BEST AVAILABLE COPY.

□ other: \_\_\_\_

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.